

Realizing Degree Imbalances in Directed Graphs

Dhruv Mubayi and Todd G. Will
Davidson College, Davidson NC 28036

Douglas B. West
University of Illinois, Urbana, IL 61801

Abstract

In a directed graph, the imbalance of a vertex is its outdegree minus its indegree. We characterize the sequences that are realizable as the sequence of imbalances of a simple directed graph. Moreover, a realization of a realizable sequence can be produced by a greedy algorithm.

A sequence of integers is *graphic* if there is a simple undirected graph for which it is the degree sequence. There are well-known characterizations of graphic sequences, both recursive and non-recursive. We obtain analogous results for the “imbalance sequences” of digraphs with no repeated arcs. We define the *imbalance* of a vertex in a digraph to be the difference between the number of exiting arcs and the number of entering arcs. Symbolically, $b(v) = d^+(v) - d^-(v)$, where $d^+(v)$ is the out-degree of v and $d^-(v)$ is the in-degree. A sequence $b = b_1, b_2, \dots, b_n$, is *realizable* if there exists a simple digraph G with vertices $v_1, v_2, \dots, v_n$ such that $b(v_i) = b_i$.

If we allow repeated arcs, then the trivial necessary condition $\sum b_i = 0$ is sufficient, using only arcs from vertices with positive imbalance to vertices with negative imbalance; this is analogous to the observation that when $\sum d_i$ is even, the sequence d of nonnegative integers is the degree sequence of some undirected graph, allowing loops and multiple edges. Hence we forbid repeated arcs. It does not matter whether we forbid loops or pairs of opposed arcs, since these have no affect on the imbalance sequence.

Our results are analogous to the known conditions for graphic sequences. Havel [2] and Hakimi [3] independently showed that a non-increasing sequence of nonnegative integers $d = d_1, d_2, \dots, d_n$ is graphic if and only if $\hat{d} = \hat{d}_2, \dots, \hat{d}_n$ is graphic, where $\hat{d}$ is formed from d by deleting d_1 and

subtracting 1 from the d_1 largest remaining elements of d . In this context, we define d_i to be *larger* than d_j if $d_i > d_j$ or if $d_i = d_j$ and $i > j$, so that $\hat{d}$ is also non-increasing. This result provides a recursive procedure that tests whether a sequence is graphic. If so, retracing the computation produces a realization; if $\hat{G}$ realizes $\hat{d}$, then adding a vertex whose neighborhood corresponds to the d_1 largest entries of $\hat{d}$ forms a graph G that realizes d .

Erdős and Gallai [1] provided a non-recursive characterization of graphic sequences: the non-negative integer sequence $d = d_1, d_2, \dots, d_n$ is graphic if and only if $\sum d_i$ is even and $\sum_{i=1}^k d_i \leq k(k-1) + \sum_{j=k+1}^n \min\{k, d_j\}$ for all $1 \leq k \leq n$. The inequalities are necessary because from the set S of k vertices with highest degree there are at most $\sum_{j=k+1}^n \min\{k, d_j\}$ edges to $V(G) - S$, and there are at most $k(k-1)/2$ edges within S . Recently, Aigner and Triesch [4] presented a short proof of sufficiency using ideals in the dominance order on sequences with fixed sum.

A greedy algorithm for realizing an imbalance sequence directs arcs from the vertex with greatest imbalance (b_1) to the b_1 vertices with smallest imbalance. This heuristic is natural in the sense that the vertices with large (positive) imbalances require high out-degree, and vertices with small (i.e. negative) imbalances require high in-degree. This suggests a modified sequence $\hat{b}$ and a recursive test, analogous to the Havel-Hakimi test.

We also prove a non-recursive characterization analogous to the Erdős-Gallai result. If $b = b_1, b_2, \dots, b_n$ is realizable, then $\sum b_i = 0$, because each edge contributes positively and negatively to the total imbalance. Also, $\sum_{i=1}^k b_i \leq k(n-k)$; letting S be the set of k vertices with largest imbalance, the edges within S contribute nothing to $\sum_{i=1}^k b_i$, and the pairs $S \times (V(G) - S)$ contribute at most one each.

These obvious necessary conditions are also sufficient; we present two proofs. The first shows that the greedy algorithm suggested above produces a realization when one exists. The second proof, based on the method of ideals used by Aigner and Triesch, is much shorter but does not yield such a fast construction algorithm.

Definition 1 *A non-increasing sequence of integers $a = a_1, a_2, \dots, a_n$ is a feasible sequence if the sum of the elements is zero and $\sum_{i=1}^k a_i \leq k(n-k)$ for all $1 \leq k \leq n$.*

From a feasible sequence $a = a_1, a_2, \dots, a_n$, we form an associated sequence $\hat{a} = \hat{a}_1, \hat{a}_2, \dots, \hat{a}_n$ by deleting a_1 and adding 1 to the a_1 smallest elements of a , where a_i is smaller than a_j if $a_i < a_j$ or if $a_i = a_j$ and $i < j$. The sequence $\hat{a}$ is the result of applying one step of the greedy algorithm to the sequence a . The complication in determining which entries of a are augmented to obtain $\hat{a}$ occurs when $a_{n-a_1+1} = a_{n-a_1}$; in this case there is a gap consisting of elements to which we add zero. We endure this complication to ensure that $\hat{a}$ is also non-increasing. The following example

with $n = 9$ and $a_1 = 5$ produces such a gap, since $a_5 = a_4 = 2$.

$$\begin{array}{cccccccccc} a : & 5 & 3 & 2 & 2 & 2 & 2 & -5 & -5 & -6 \\ & & \cdot & 0 & 1 & 1 & 0 & 0 & 1 & 1 & 1 \\ \hat{a} : & \cdot & 3 & 3 & 3 & 2 & 2 & -4 & -4 & -5 \end{array}$$

The values of a_i in the gap and the values of a_i to the left of the gap are all equal. When this occurs in a feasible sequence, a stronger statement can be made about the partial sums.

Lemma 2 *If a is a feasible sequence and $a_k = a_{k+1} = \cdots = a_{k+m}$, then $\sum_{i=1}^k a_i \leq k(n-k) - m$.*

Proof: If $a_k \leq n - 2k - m$, then

$$\sum_{i=1}^k a_i = \left(\sum_{i=1}^{k-1} a_i \right) + a_k \leq (k-1)(n-k+1) + (n-2k-m) = k(n-k) - m - 1.$$

If $a_k > n - 2k - m$, then

$$\sum_{i=1}^k a_i = \left(\sum_{i=1}^{k+m} a_i \right) - m a_k \leq (k+m)(n-k-m) - m(n-2k-m+1) = k(n-k) - m. \quad \square$$

Theorem 3 *If a is feasible, then $\hat{a}$ is feasible.*

Proof: Because $\sum_{i=1}^n a_i = 0$ and $a_1 \leq 1 \cdot (n-1) = n-1$, the sequence $\hat{a}$ is well defined. By the definition of the a_1 “smallest” elements, $\hat{a}$ is non-increasing. The construction of $\hat{a}$ distributes a_1 among the other entries, so the sum is still 0. It remains to verify the condition on partial sums, which is $\sum_{i=2}^k \hat{a}_i \leq (k-1)(n-k)$ for $2 \leq k \leq n$.

Suppose $a_1 = n - r$, and suppose the construction of $\hat{a}$ involves a gap of s entries to which we add 0, where $s \geq 0$. Beginning with the leftmost entry to which 1 is added, there are $(n-r) + s$ positions in the sequence; hence $s < r$ and a_{r-s+1} is the leftmost entry to which 1 is added. If $k \leq r - s$, then

$$\sum_{i=2}^k \hat{a}_i = \sum_{i=2}^k a_i \leq (k-1)(n-r) \leq (k-1)(n-k).$$

Now suppose $k > r - s$. Let $\alpha = \sum_{i=2}^k \hat{a}_i - \sum_{i=2}^k a_i$; this is the number of added 1's in the sum. Let t be the number of positions in the sequence that are to the left of the gap and receive an augmentation of 1 ($t = 0$ if $s = 0$). The rightmost entry to which 0 is added is a_{r+t} .

$$\begin{array}{cccccccccccccccc} & & & & & \overbrace{\hspace{2cm}}^t & & \overbrace{\hspace{2cm}}^s & & & & & & & & \\ a : & a_1 & a_2 & \cdots & a_{r-s} & a_{r-s+1} & \cdots & a_{r-s+t} & a_{r-s+t+1} & \cdots & a_{r+t} & a_{r+t+1} & \cdots & a_n \\ & & 0 & \cdots & 0 & 1 & \cdots & 1 & 0 & \cdots & 0 & 1 & \cdots & 1 \\ \hat{a} : & \hat{a}_1 & \hat{a}_2 & \cdots & \hat{a}_{r-s} & \hat{a}_{r-s+1} & \cdots & \hat{a}_{r-s+t} & \hat{a}_{r-s+t+1} & \cdots & \hat{a}_{r+t} & \hat{a}_{r+t+1} & \cdots & \hat{a}_n \end{array}$$

If $r - s < k < r + t$, then $a_k = \dots = a_{r+t}$. By the Lemma, $\sum_{i=1}^k a_i \leq k(n - k) - (r + t - k)$. In this case, $\alpha \leq t$. If $k \geq r + t$, then $\sum_{i=1}^k a_i \leq k(n - k)$ and $\alpha = k - r$. In either case, we have $(\sum_{i=1}^k a_i) + \alpha \leq k(n - k) + (k - r)$. Hence

$$\sum_{i=1}^k \hat{a}_i = \left(\sum_{i=1}^k a_i \right) + \alpha - a_1 \leq k(n - k) + (k - r) - (n - r) = (k - 1)(n - k). \quad \square$$

A simple inductive argument now completes the proof of the sufficiency of the condition and shows that the greedy algorithm produces a realization if one exists. We also present a second proof, using the sufficiency method of Aigner and Triesch [4]. The idea here is to define a partial order P on sequences such that 1) the realizable sequences form an ideal (down-set), and 2) the maximal sequences in P that satisfy the desired condition (here feasibility) are realizable. This implies that all sequences satisfying the condition are realizable, because they are dominated in P by a realizable sequence.

Corollary 4 *A sequence a is realizable as an imbalance sequence if and only if a is feasible.*

Proof: 1: We have already argued the necessity of the condition. For sufficiency, we used induction on the length n of a . If $n = 1$, then $a_1 = 0$ and K_1 has imbalance sequence a . If $n > 1$, then Theorem 3 implies that $\hat{a}$ is a feasible sequence of length $n - 1$. By the induction hypothesis, there is a graph G' with vertices $v_2, v_3, \dots, v_n$ such that $b(v_i) = \hat{a}_i$. Form G by adding the vertex v_1 and arcs from v_1 to v_j for each $j \in V(G')$ such that $\hat{a}_j = a_j + 1$. The graph G realizes a .

Proof: 2: Consider the domination order P on non-increasing sequences of integers with sum 0; the order is defined by $a \geq b$ if $(\sum_{i=1}^k a_i) \geq (\sum_{i=1}^k b_i)$ for all k . In the subposet of P induced by the feasible sequences, we claim that only the sequence $n - 1, n - 3, \dots, -(n - 3), -(n - 1)$ is maximal. It achieves each constraint with equality, and therefore by the definition of the order it dominates all other feasible sequences. Furthermore, this is the imbalance sequence of the transitive tournament with n vertices and hence is realizable.

It remains to show that the realizable sequences in P form an ideal. If $b > b'$ in P , then b' is obtained from b by a sequence of unit shifts in which some position i decreases by 1 and some later position j increases by one; it suffices to show that every such shift maintains realizability. Suppose b is realizable, and consider positions i, j for such a shift. Because $b_i > b_j$, there is some other vertex z in the realization such that $i \rightarrow z \not\rightarrow j$ or $i \rightarrow z \rightarrow j$ or $i \not\rightarrow z \rightarrow j$. In these three cases, respectively, we transform the graph to $i \not\rightarrow z \leftarrow j$ or $i \not\rightarrow z \not\rightarrow j$ or $i \leftarrow z \not\rightarrow j$. The only change in the sequence is that b_i goes down by 1 and b_j goes up by 1, so we have realized b' also as an imbalance sequence. $\square$

References

- [1] P. Erdős and T. Gallai, Graphs with prescribed degrees of vertices, *Mat. Lapok* **11** (1960), 264–274 (in Hungarian).
- [2] V. Havel, A remark on the existence of finite graphs, *Casopis Pest. Mat.* **80** (1955), 477–480.
- [3] S. L. Hakimi, On the realizability of a set of integers as degrees of the vertices of a graph, *SIAM J. Appl. Math.* **10** (1962), 496–506.
- [4] M. Aigner and E. Triesch, Realizability and uniqueness in graphs, *Discrete Math.* **136** (1994), 3–20.